

RR LYRAE – THEORY vs OBSERVATION

ZOLTÁN KOLLÁTH¹, J. ROBERT BUCHLER² & MICHAEL FEUCHTINGER²

ASTROPHYSICAL JOURNAL, revised version, in press

ABSTRACT

The luminosities, effective temperatures and metallicities that are derived empirically by Kovács and Jurcsik from the light curves of a large number of globular cluster and field RRab and RRc stars are compared to theoretical RR Lyrae models. The strong luminosity dependence of the empirical blue and red edges ($\text{Log } L$ vs. $\text{Log } T_{\text{eff}}$ diagram) is in disagreement with that of both radiative and convective models. A reexamination of the theoretical uncertainties in the modelling leads us to conclude that the disagreement appears irreconcilable.

Subject headings: stars: variable, stars: oscillations, stars: RR Lyrae, stars: horizontal-branch, stars: abundances, stars: distances, stars: atmospheres, stars: fundamental parameters, globular cluster

¹Konkoly Observatory, Budapest, HUNGARY; kollath@konkoly.hu

²Physics Department, University of Florida, Gainesville, FL, USA; buchler@phys.ufl.edu, fm@phys.ufl.edu

1. INTRODUCTION

The recent work of Kovács & Jurcsik (Kovács & Jurcsik 1996 [KJ96], 1997 [KJ97], Jurcsik 1998 [J98]) proposes an almost purely empirical method of extracting the absolute magnitudes, colors and effective temperatures (T_{eff}) and metallicities directly from the observed periods and light curves of RR Lyrae stars. The only theoretical input appears in the transformation from M_V and $V - K$ to L and T_{eff} via Kurucz's static model atmospheres. The very empirical nature and potential usefulness of the approach has attracted a great deal of attention from observers.

The most recent work (J98) compiles and analyzes a sizeable set of observational data of RR Lyrae cluster variables and RR Lyrae field stars. This study thus includes RR Lyrae stars with metallicities ranging from $Z=0.00001$ to almost solar $Z=0.020$. The end products of her analysis that we are most concerned about here are $\text{Log } L$ - $\text{Log } T_{\text{eff}}$ and $\text{Log } L$ - $\text{Log } P$ plots, and the byproducts which are relations between luminosity L , mass M and metallicity Z .

The J98 $\text{Log } L$ - $\text{Log } T_{\text{eff}}$ and $\text{Log } L$ - $\text{Log } P$ data are shown in Figure 1 as small dots, circles for RRab and triangles for RRc stars. The figures indicate well defined slopes for the fundamental and overtone blue edges and red edges, all four of which have approximately the same values, although there may be a slight broadening of the instability strip with luminosity. Hereafter we call this slope Ξ ($= \Delta \text{Log } L / \Delta \text{Log } T_{\text{eff}}$).

J98 pointed out that no model calculations can explain the strong dependence of the temperature on the luminosity. In §2 we first confirm that indeed the slope Ξ of this empirical $\text{Log } L$ vs. $\text{Log } T_{\text{eff}}$ diagram is in irreconcilable disagreement with that of radiative models. Next, in §3 we show that the inclusion of turbulent convection in the models only shifts the $\text{Log } L$ vs. $\text{Log } T_{\text{eff}}$ line, but does not change the slope. The discrepancy is therefore very basic because it is already inherent in models that are as fundamental as purely radiative ones. In §4 we reexamine the uncertainties of both radiative and turbulent model

calculations and conclude that the theoretical Ξ is very robust and cannot be changed very much without introducing new physics. In §5 we discuss the theory that goes into the empirical relations of Jurcsik, in particular the influence of the use of the static Kurucz color – temperature transformation. We conclude in §6.

2. RADIATIVE MODELS OF RR LYRAE STARS

In the following we examine how well RR Lyrae models agree with the J98 data. We first examine radiative models, and thus limit ourselves to the vicinity of the blue edges.

Figure 1 shows a $\text{Log } L$ - $\text{Log } T_{\text{eff}}$ plot on the left. On these we superpose the linear fundamental and first overtone blue edges of radiative RR Lyrae models. We recall here that among the linear edges of the instability strip only the overtone linear blue edge and the fundamental linear red edge coincide with the observable edges of the instability strip. Nonlinear dynamical effects shift the observable fundamental blue edge and the overtone red edge to the red and to the blue, respectively, compared to their corresponding linear edges, by up to several 100 K (*e.g.*, *cf.* Buchler 2000). Strictly speaking, for radiative models, it is only the linear overtone blue edge that is relevant, but the linear fundamental blue edge can give a rough indication of the slope Ξ . Furthermore, the red edges are determined by convection, and radiative red edges are not relevant and are therefore not shown in Fig. 1.

The solid lines have been computed for models with $M=0.65$, $X=0.75$ and $Z=0.001$. The dotted lines represent models with M and Z from Jurcsik's $\{L, M, Z\}$ relations, the dashed lines those with M from Jurcsik, but with $Z=0.001$. Finally, the large open circles are models from Tuggle and Iben (1972) for $M=0.6$ and $Z=0.001$ (with the old Los Alamos opacities). They have essentially the same Ξ as ours which are computed with the OPAL opacities (Iglesias & Rogers 1996) merged with the low temperature Alexander-Ferguson (1994) opacities.




Fig. 1 : Left: $\text{Log } L - \text{Log } T_{\text{eff}}$ plot: RRab (filled circles) and RRc (triangles) stars from Jurcsik ([J98] and priv. comm.); Theoretical fundamental and first overtone blue edges for radiative models: *solid lines*: fixed mass and Z ; *dotted lines* have M and Z from Jurcsik's relation; *dashed lines* have M from Jurcsik with fixed $Z=0.001$; Tugge & Iben's overtone and fundamental blue edges are reported as open circles and squares, respectively. Right: Same data in a $\text{Log } L - \text{Log } P$ plot.

Again, because we are computing radiative models we do not expect the location of the blue edge to be in perfect agreement with the edge of the instability strip, but we note a very large discrepancy in the slope Ξ .

Figure 1 also displays a $\text{Log } L - \text{Log } P$ plot on the right. Only the slope of the constant mass models is in almost acceptable agreement with the J98 data, but those calculated with the J98 $\{M, L, Z\}$ are in strong disagreement.

In Figures 2 we display again a $\text{Log } L - \text{Log } T_{\text{eff}}$ plot which shows the effect of composition on the location of the linear blue edges. The (radiative) models were computed with $M=0.65$. Here the solid lines have $X=0.75$, $Z=0.001$, the dotted lines have $X=0.75$, $Z=0.004$ and the dashed lines have $X=0.70$, $Z=0.001$.

The location of the blue edge displays very little sensitivity to either metallicity Z or helium content Y , within a reasonable range of values. Adjusting the helium content or metallicity does not provide a resolution of the slope discrepancy. We describe further tests in §4.

We conclude this section by noting that purely radiative RR Lyrae models are in severe disagreement with Jurcsik's $\text{Log } L - \text{Log } T_{\text{eff}}$ data.

3. CONVECTIVE MODELS OF RR LYRAE STARS

The addition of time-dependent turbulent convection in the models has led to some spectacular successes compared to radiative models, in particular, in finally predicting double mode behavior both in RR Lyrae (Feuchtinger 1998) and in Cepheids (Kolláth, Beaulieu, Buchler & Yecko 1998). One would hope therefore that the inclusion of turbulent convection might also remove the discrepancy between observations and theory described in the previous section.

Yet, in that respect all of our modelling efforts with turbulent convection have proved in vain, despite the flexibility afforded by the 8 free order unity (α) parameters that the turbulent convective equations contain (*e.g.*, Yecko, Kolláth & Buchler 1997, Buchler & Kolláth 2000). We find that turbulent convection can shift the blue edges, but cannot produce the differential effect with respect to luminosity that is required to give the right slope Ξ in the $\text{Log } L$ vs. $\text{Log } T_{\text{eff}}$ plots. In Figure 3 we show the results for three different combinations of these parameters.

We note that different codes and slightly different recipes for convection give essentially the same theoretical slopes. For example, even though no linear models are computed, the nonlinear hydrodynamical models of Bono *et al.* (1997) indicate a similarly steeper slope than that of J98.

4. DISCUSSION OF THE MODELING ASSUMPTIONS

In this section we discuss the effects of several of the approximations that are inherent in the numerical modeling to see how robust the theoretical $\text{Log } L - \text{Log } T_{\text{eff}}$ slope is.

4.1. Equation of State

Our code uses as equation of state a simple iteration of the Saha equations (*e.g.* Stellingwerf 1982). This equation of state is very similar to the OPAL (Rogers *et al.* 1996) and the MHD (Däppen *et al.* 1988) equations of state and we do not believe that the tiny differences can be responsible for the discrepancy between the models and the J98 data.

4.2. Radiative Transport

Our radiative code uses a standard equilibrium diffusion approximation for the radiative transport, *i.e.*,

$$L = (4\pi R^2)^2 \frac{c}{3\kappa} \frac{\partial}{\partial m} aT^4 \quad (1)$$

in which one uses for the opacity κ the Rosseland mean (*e.g.*, Mihalas & Mihalas 1984). Eq. 1 has several shortcomings in the region above the photosphere. First, it implies an Eddington factor of $f_E = 1/3$, which is not correct in the optically thin outer region where f_E approaches 0.4–0.5 (Feuchtinger & Dorfi 1994, Fig. 3). Second, in this regime, the opacities should be higher than those given by the Rosseland mean (Alexander & Ferguson 1994). Finally, for the computation of the periods and growth rates it would be more appropriate to linearize the radiation hydrodynamics equations rather than the equilibrium diffusion equation. The magnitude of the errors introduced by the first effect can easily be estimated.

One can approximate the effect of the Eddington factor in Eq. 1. by replacing 3κ by κ/f_E . The Eddington factor also appears in the radiation pressure which becomes $p_{\text{rad}} \rightarrow 3f_E p_{\text{rad}}$. We can disregard sphericity effects in the momentum equation contained in the term $(p_{\text{rad}} - f_E e_{\text{rad}})/r$.

To simulate the effect of an increasing f_E we assume that for $T < T_{\text{eff}}$, f_E increases smoothly from 1/3 to 1/2 with decreasing temperature. The results confirm our hunch that an increase of f_E in the very outer region has very little influence on the growth rates and almost none on the periods. More importantly it has no differential sensitivity to luminosity.




Fig. 2 : Log L -Log T_{eff} plot: RRab (filled circles) and RRc (triangles) stars: Convective models: Solid, dotted and dashed lines are for three different combinations of the α parameters in the convective model equations, chosen to give a reasonable width for the instability strip.




Fig. 3 : Log L -Log T_{eff} plot: RRab (filled circles) and RRc (triangles) stars: Convective models: Solid, dotted and dashed lines are for three different combinations of the α parameters in the convective model equations, chosen to give a reasonable width for the instability strip.

We have artificially, and somewhat arbitrarily, increased the opacity from the vicinity of T_{eff} outward. We have not found an appreciable differential effect with T_{eff} , and have therefore not pursued this avenue with more detailed modelling.

Finally, the insensitivity of the slope to f_E makes us believe that the linearization of the radiation hydrodynamics equations instead of the common equilibrium diffusion equation (requiring a serious coding effort) would also make little difference.

We conclude that the discrepancy between the models and the observations is not due to an inadequate treatment of the radiation transport.

4.3. Composition

Could compositional make-up be more important than expected? We have computed a number of RR Lyrae models with various combinations of Y and Z, but find that the effect on Ξ is negligible. Next we have artificially increased the abundances of the easily ionizable light elements such as Mg and Na in the OPAL opacities. Again this has such a negligible effect on Ξ that the inclusion of this data would unnecessarily clutter Figure 2.

An inhomogeneous composition is unlikely to exist in RR Lyrae envelopes because of convection, but even if existed our tests with various changes in composition make us doubt that it would resolve the discrepancy.

4.4. Rotation

Could rotation be responsible for the discrepancy? In order to estimate the magnitude of the effect of rotation we have included a spherical pseudo-centrifugal acceleration $\omega^2 r$ in the equilibrium model and in the computation of




Fig. 4 : Left: T_{eff} -(V-I) plot and right: (V-I)- M_V plot: Theoretical color variations for two nonlinear full amplitude RRab models with different luminosities of $L = 40 L_{\odot}$ (dashed line) and $L = 52 L_{\odot}$ (full line). Crosses indicate the time steps of the nonlinear calculations. See text for details.

the periods and growth rates. We find that a rather short rotation period, of order of a few days, would be necessary to have an impact. Furthermore the disagreement is worsened by rotation because the slope steepens.

We conclude that moderate rotation rates cannot be the cause of the discrepancy.

4.5. Evolutionary Effects

The blueward moving evolutionary paths turn around at some T_{eff} and then move upward (*e.g.*, Dorman 1992, Lee, Demarque & Zinn 1990). If this happened inside the linear instability strip, then the leftmost blue edge, that of the overtone mode, would be defined by the topology of these paths, rather than by the pulsational stability of the models. However, the fact that the Jurcsik blue and red edges of the fundamental and the overtone are all essentially parallel eliminates this possibility as an explanation for the shallow observed slope. Moreover, for some of the clusters, full photometry of the horizontal branch exists, and there is no indication of any gap next to the RR Lyrae region. Clearly there are stars next to the empirical red edge for which the amplitude (if not zero) is less than the observational limit.

4.6. Conclusion

The theoretical Ξ slope is very robust with respect to the uncertainties or approximations inherent in our code, but is in disagreement with the Jurcsik empirical relations. We have therefore been led to reexamine the latter for their robustness, especially since they also make use of some theory, *viz.* color to temperature transformation.

5. DISCUSSION OF THE EMPIRICAL RELATIONS

Because of the strong disagreement between Jurcsik's processed observational data and the predictions from something as basic as radiative models, it is worth while to reexamine some of the most uncertain points in the reduction of the observational data.

The observational material consists of RRab periods and lightcurves. KJ96, JK97 and J98 deduce the color, the visual magnitude M_v and the metallicity Z ($[\text{Fe}/\text{H}]$) in the form of linear fully empirical relations

$$\begin{pmatrix} M_v \\ V - K \\ [\text{Fe}/\text{H}] \end{pmatrix} \equiv c_0 + A \cdot \begin{pmatrix} P \\ A_1 \\ \phi_{31} \\ \phi_{41} \end{pmatrix} \quad (2)$$

The connection with $\text{Log } T_{\text{eff}}$ and with $\text{Log } L$ was then made through static envelope models of Kurucz (1993, *cf.* J98 for details). This is the only theoretical input into the otherwise empirical relations.

$$\begin{pmatrix} \text{Log } L \\ \text{Log } T_{\text{eff}} \end{pmatrix} = c_1 + C \cdot \begin{pmatrix} M_v \\ V - K \\ [\text{Fe}/\text{H}] \end{pmatrix} = c_2 + C \cdot A \cdot \begin{pmatrix} P \\ A_1 \\ \phi_{31} \\ \phi_{41} \end{pmatrix} \quad (3)$$

5.1. Color - T_{eff} Transformation

J98 uses the *static* envelope calculations of Kurucz's ATLAS code (for details *cf.* J98), to make the transformation from average color to average T_{eff} . The T_{eff} however




Fig. 5 : Left: R Rab Period – Radius plot; *open circles*: Baade-Wesselink radii, and *crosses*: same stars from J98; The solid and dashed lines are linear regressions through the Baade-Wesselink and the Jurcsik data, respectively. Right: *thick lines*: radiative models with Jurcsik L, M, Z relation; *thin lines*: radiative models with constant mass and luminosity; *filled squares*: same stars from J98;

varies considerably during one pulsation cycle, compared to the width of the instability strip, and the question arises whether the correct mean color and mean temperature are still related through the static Kurucz relations. To check the validity of this assumption we have computed the spectra of two full amplitude RR Lyrae models.

We investigate the theoretical color variation for two RR Lyrae models with different luminosities ($L = 40 L_{\odot}$ and $L = 52 L_{\odot}$, both for $M = 0.65 M_{\odot}$, $T_{\text{eff}} = 6500 \text{ K}$, $X = 0.76$ and $Z = 0.001$). Nonlinear full amplitude models are taken from Feuchtinger (1999) and the color variations in V and I are computed as outlined in Dorfi & Feuchtinger (1999). The results can be inferred from Fig. 4.2 which displays the color difference $V - I$ as a function of the effective temperature (left panel) and the corresponding color-magnitude diagram (right panel), both over one oscillation cycle. Full and dotted lines refer to the $L = 40 L_{\odot}$ and $L = 52 L_{\odot}$ models, respectively. The differential effect with L is seen to be small. Dynamical atmospheric effects thus cannot account for the discrepancy.

There remains the question of the consistency of the static Kurucz atmospheres with our frequency dependent calculation. For that purpose we have checked the color – temperature transformation against the Kurucz tables at selected points on the pulsation cycle. The agreement is quite satisfactory, and any discrepancies are irrelevant for our purpose here.

5.2. RR Lyrae Period - Radius Relation

The radii of RR Lyrae stars can be obtained independently either from a Baade-Wesselink approach or from the J98 empirical L and T_{eff} . In Fig. 5, on the left, we have plotted as open circles the Baade-Wesselink radii of the 15 RRab stars of Jones *et al.* (1992) (we have omitted RR Leo). Superposed as crosses are the radii for the same stars derived from the J98 L and T_{eff} . The agreement is remarkably good, but worsens for the longer periods. The solid and dashed lines represent linear regression lines for the BW and for the J98 period – radius relations.

We have also derived the radii of the additional J98 stars that were not in the Jones sample and show them as small dots in the righthand figure. The thin lines represent constant mass and constant luminosity radiative models, and the thick lines radiative models that obey the J98 L, M, Z relations. Since the radiative models do not provide a red edge, the upward extent of the thin or thick lines is not significant. The latter models are seen to give better agreement with the swarm of the J98 empirical radii.

We conclude that the agreement between the empirical J98 and Jones' Baade-Wesselink radii is remarkably good, and so is the agreement that can be achieved with theoretical models. A consideration of the period – luminosity relations therefore does not give us any clues as to the origin of the slope discrepancy.

6. CONCLUSIONS

We have shown that the Jurcsik relations lead to RR Lyrae radii that are in good agreement with both Baade-Wesselink radii and with theoretical radii. On the other hand, there exists a strong discrepancy between the slope of the theoretical $\text{Log } L - \text{Log } T_{\text{eff}}$ relation and the slope of the empirical Jurcsik relation.

We have reviewed the physical and numerical uncertainties that enter the theoretical calculations. The discrepancy exists already at the level of purely radiative modelling. We have further shown that our model equations for turbulent convection do not alter the slope of the $\text{Log } L - \text{Log } T_{\text{eff}}$ relation despite the large number (8) of adjustable parameters (α 's) that we have at our disposal. Consequently, apart from the uncertainties inherent in a 1D recipe for turbulent convection, it appears that the discrepancy is not caused by a deficiency of the theoretical models. We note that the same slope discrepancy is implicit in older and more recent independent calculations (Tuggle & Iben 1972, Bono *et al.* 1997).

The derivation of the empirical relations makes use of a *static* Kurucz color - T_{eff} transformation. By computing the behavior of color versus temperature over the pulsation cycle we have shown that the use of static envelopes is in fact a very good approximation, and therefore cannot be the culprit.

Finally, the shape of the evolutionary tracks, through their potential avoidance of certain regions of the instability strip cannot be responsible either.

At this time the origin of this disturbing discrepancy constitutes an unsolved puzzle. It remains to be seen whether an improved treatment of turbulent convection or more complete observations will resolve the difficulty.

7. ACKNOWLEDGEMENTS

The authors would like to thank Johanna Jurcsik for kindly providing us with copies of her data sets. They also acknowledge fruitful discussions with Johanna Jurcsik and Géza Kovács. This work has been supported by NSF (AST 95-28338 and AST 98-19608) and the Hungarian OTKA (T-026031).

REFERENCES

- Alexander, D. R. & Ferguson, J. W. 1994, *ApJ* 437, 879
 Bono, G., Caputo, F., Castellani, V. & Marconi, M. 1997, *A&ASuppl* 121, 327
 Buchler, J. R. 2000, in *The Impact of Large-Scale Surveys on Stellar Pulsations*, Eds. L. Szabados & D. Kurtz, ASP Conf. Ser. (in press)
 Buchler, J. R. & Kolláth, Z., 2000, *Turbulent Convection in the Classical Variable Stars*, in *Astrophysical Turbulence and Convection*, Eds. J.R. Buchler & H. Kandrú, Annals of the New York Academy of Sciences, 898, 39.
 Cohen, J. G. 1992, *ApJ* 400, 528
 Däppen, W., Mihalas, D., Hummer, D. G., Mihalas, B. W., 1988, *ApJ* 332, 261
 Dorman, B. 1992, *AJ Suppl Ser.* 81, 221
 Dorfi, E. A. & Feuchtinger, M. U. 1999, *AA* 348, 815
 Feuchtinger, M. U. 1998, *AA* 337, L29
 Feuchtinger, M. U. 1999, *AA* 351, 103
 Feuchtinger, M. U. & Dorfi, E. A. 1994, *AA* 291, 209
 Iglesias, C.A. & Rogers, F. J., 1996, *ApJ*, 464, 943
 Jones, R. V., Carney, B. W., Storm, J. & Latham, D.W. 1992, *ApJ* 386, 646
 Jurcsik, J., 1998, *AA* 333, 571
 Kovács, G. & Jurcsik, J. 1996, *ApJLett* 466, L17
 Kovács, G. & Jurcsik, J. 1997, *AA* 322, 218
 Kolláth, Z., Beaulieu, J.P., Buchler, J. R. & Yecko, P., 1998, *ApJ Lett* 502, L55
 Lee, Y. W., Demarque, P. & Zinn, R. 1990, *ApJ* 350, 155
 Liu, T. & Janes, U. A. 1990, *ApJ* 354, 273
 Mihalas, D. & Mihalas, B. 1984, *Foundations of Radiation Hydrodynamics*, (Oxford: University Press)
 Rogers, F.J., Swenson, F.J. and Iglesias, C. A., 1996, *ApJ* 456, 902
 Stellingwerf, R.F., 1982, *ApJ* 262, 330
 Tuggle, R. S. & Iben, I. 1972, *ApJ* 178, 455
 Yecko, P., Kolláth, Z. & Buchler, J. R. 1998, *Turbulent Convective Cepheid Models: Linear Properties*, *Astronomy & Astrophysics* 336, 553–564